



revolute; *column* stout, semiterete, 1.5 mm long, the anther and bilobed stigma apical.

Etymology: Named for Bryon Rinke of Winfield, KS, who cultivates this species

Among the species of section *Macrophyllae-fasciculatae*, this species is distinguished by a pendent, narrowly ovate leaf, a colorful flower subtended by a reclining spathe, and a thick, convex, panduriform lip.

Acronia rinkei Luer, *sp. nov.* TYPE: ECUADOR. without collection data, purchased from Ecuagenera, flowered in cultivation in Winfield, Kansas, September 2010, by *Bryon Rinke 0701* (Holotype: MO), C. Luer illustr. 21562. Fig. 6.

Inter species Aconitiae Presl sect. Macrophyllae-fasciculatae (Lindl.) Luer, *folio anguste cordato dependens, spatha decumbenti, sepalo dorsali synsepaloque obtusis, petalis anguste acutis, et labello panduriformi lateribus supra basim involutis pubescentibus distinguitur.*

Plant medium in size, epiphytic, caespitose; roots slender. *Ramicauls* slender, erect, 14 cm long, with a close, tubular sheath below the middle. *Leaf* pendent, coriaceous, narrowly ovate, acute, 9.5 cm long, 3 cm wide, the base cordate, sessile. *Inflorescence* a single flower, borne from a reclining spathe 6–7 mm long at the base of the leaf; peduncle 3–5 mm long, within the spathe; floral bract 5–6 mm long; pedicel 5–6 mm long; ovary 4 mm long; *sepals* fleshy, glabrous, the dorsal sepal erect, yellow, oblong, minutely apiculate at the rounded apex, 13 mm long, 5 mm wide, 3-veined, the lateral sepals red-purple, connate to the tips into an ovate, obtuse synsepal, 12 mm long, 7.5 mm wide, 6-veined; *petals* yellow, narrowly ovate, acute, 7 mm long, 1.25 mm wide, 1-veined; *lip* red-purple, fleshy, panduriform, pubescent, convex, 6 mm long, 3 mm wide at the base, 1 mm wide near the middle, the apex obtuse to rounded, the base with a small glenion, the sides