

BIPINNULA.

Juss. gen. 65.

CHAR. GEN. Pollinia duo pulvrea bipartita, caudiculâ glandulâque nullis.
Anthera terminalis opercularis bilocularis . . . Stigma
Columna semiteres membranaceo-marginata. Labellum anticum cucullatum
integrum cristatum appendicibus filiformibus fimbriatum. Sepala ringentia
membranacea; supremis nanis; inferioribus labello suppositis, anamorphose
solutâ, longissimis fimbriatis. *Herbæ terrestres* (Americæ Australis tempe-
ratae) radicibus fasciculatis. Folia plana radicalia. Scapi squamosi. Flores
colorati?

1. *B. plumosa*; scapo folioso multifloro, sepalis inferioribus
labelloque plumosis.

In pascuis herbidis ad vicum *Colina* urbis *Santiago* finitimum semel vidit
McRae, Novembre floridam. (v. s. sp.)

Stem about a foot and a half high. *Leaves* erect, oblong-lanceolate,
somewhat obtuse, gradually changing to long, membranous, acuminate,
sheathing squamæ. *Spike* about five inches long, eight or ten
flowered. *Bractæ* membranous, acuminate, nearly as long as the
upper sepals. *Flowers* ringent. *Upper sepal* lanceolate acute, *inner*
erect, rather shorter, obtuse; *lower sepals* three times as long as the
upper ones, ovate-acuminate, spreading, membranous, finely fringed
beyond the middle with long delicate filiform processes which arise
from the margin. *Labellum* ovate-acuminate, cucullate at base, about
one-third the length of the lower sepals, like which it is also bordered
with a fringe of long filiform ciliæ. *Columna* erect, half the length
of the upper sepals.

Two specimens only of this noble species, one in flower, the
other in bud, were discovered by Mr. M^eRae in the grassy
pastures about the hot-baths of Colina, a small village on the
skirts of the plain which surrounds the town of Santiago. The
fine fringe that borders the lower sepals and the labellum call
to mind some of the North American species of *Habenaria*.
The flowers measure, from the base of the ovary to the apex
of the lower sepals, fully three inches.