

116. — h. — ONCIDIUM GIREOUDIANUM Rehb. fl. Mss. Huc: *Brassia Gireoudiana*
Rehb. fl. Warscewicz, Rehb. fl. Gtz. — Xenia p. 79:

Aff. *Brassiae guttatae* Lindl. sepalis labello ter — quater longioribus, labello a basi latoligulata utrinque obtusangula subito dilatato, hac parte triangula anteriori basi utrinque rotundata, lamellis quintae labelli aequilongis carinaeformibus velutinis, basi obtusangulis, tum humilioribus, dein acutangulis, demum semiovatis.

? Sub: *Brassia maculata* R. Br. *B. guttata*: in a specimen found by Mr. Skinner at San Salvador the flowers are twice as large" Lindl. Fol. V. *Brassia* p. 2.

Brassia Gireoudiana Rehb. fl. Wzwy. in Otto und Dietrich Allg. Gartenztg XXII. 1854. Nr. 35. p. 273.

Pseudobulbus prope spithamaeus, oblongo anceps, articulus superior Brassiarum more nunc manifeste evolutus. Folia a cuneata basi oblonga, obtuse acuta. Pedunculus usque bipedalis et tredecimflorus. Flores infimi ab apice supremi sepalis ad apices sepalorum lateralium septem — octopollicares. Sepala linearia longe caudata, lateralia dorsali longiora flavoviridia, demum flava fasciis paucis nunc interruptis purpureo atroviolaceis in basi. Tepala dimidio sepalo dorsali aequalia linearia acuminata dimidio inferiori purpurea atroviolacea. Labellum basi oblonga subito cordato oblongove dilatatum acutiuseculum ejusdem coloris, maculis quibusdam violaceo atropurpureis. Carinae basilares bipartitae: laminae posticae basilares a basi altiori descendentes, hinc obtuse triangulae, intus aurantiaco velutinae. Gynostemium basin versus magis antrorsum proclive, dum in *B. guttata* Lindl. rectum.

Diese prächtige Art, welche an Grösse den schönsten Formen der *B. brachiata* Lindl. gleichkommt, entdeckte Herr von Warscewicz in Costa Rica.