## TWO NEW ORCHIDS FROM JAMAICA.

BY WILLIAM FAWCETT, B.Sc., F.L.S.

**Epidendrum tridentatum**, sp. n. Stem equal at base, leafy to inflorescence. Leaves alternate, oblong or oblong-lanceolate, acute, clasping. Inflorescence a raceme or panicle, terminal, nodding; sterile bract none or one, floral bracts awl-shaped. Flowers greenishyellow. Sepals spathulate-lanceolate or broadly lanceolate, acute. Petals shorter than sepals, spathulate-linear. Labellum adnate to column, 3-6-lobed, 2-callous at base; lateral lobes trapezoidroundish, middle smaller, of 3 triangular lobules.

Habitat. Blue Mountains, Jamaica. Coll. W. Harris, April, 1893.

Plant  $1\frac{1}{2}-2$  ft. high. Leaf 5-7 in. long, 10-12 lines broad. Sterile bract 6 lines long. Floral bracts 3-5 lines long. Sepals 8-9 lines long.

Pleurothallis uncinata, sp. n. Stem biangular, about as long as the leaf, with appressed flaccid sheaths at the nodes. Leaf lanceolate, 5 or 6 times as long as the peduncles (8-9 in. long, nearly 2 in. broad). Peduncles 1-flowered, solitary or fascicled (2 or 3). Sepals oblong-lanceolate, lateral shortly connate at base, sometimes cohering above, externally scabrous, purplish, 7 lines long. Petals linear-lanceolate, half as long as sepals, orangecoloured at base, purplish above. Labellum as long as petals, indistinctly 3-lobed; lateral lobes almost obsolete, but furnished each with a hook-like process embracing the column; middle lobe oblong, blunt, minutely fimbriate-serrate, purplish. Habitat. Near Newcastle, Jamaica. Coll. W. Harris, June, 1893.