

130. *CHŸŠĪS lævis*; bracteis brevibus ovatis pedicelli longitudine, sepalo dorsali lineari-oblongo lateralibus acuminatis, petalis falcatis, labelli lobis lateralibus falcatis apice rotundatis supra columnam convergentibus intermedio membranaceo crispo subrotundo emarginato lamellis 5 carnosis glaberrimis parallelis lateralibus minoribus, columnâ basi altè excavatâ.

Of this singular Orchidaceous genus two new species have appeared in the collection of Mr. Barker. The present has very large yellow flowers, whose lip is spotted with dull brown, and more membranous than in the others; and it is remarkable for its column becoming hard and very tumid after fertilization has occurred, and eventually acquiring a green colour. The other species has white flowers, with a deep yellow fleshy lip, which is divided into two lobes, each folded twice, so as to give the whole the appearance of being four-lobed. It may be named and defined as follows :

131. *CHŸŠĪS bractescens*; bracteis cucullatis venosis foliaceis ovario longioribus, sepalis petalisque ovatis obtusis, labelli lobis lateralibus obtusis intermedio carnosio bilobo plicato lamellis 5 carnosis subæqualibus parallelis basi pubescentibus, columnâ latissimâ carnosâ cymbiformi anticè pubescente.