2a. Dichaea fragrantissima Folsom, Orchid Digest 58: 99. 1994. TYPE: Panama. Coclé, Alto del Calvario, cont. divide, N of El Copé, 850-950 m, flowered in cult., 6 Apr. 1985, J. P. Folsom 11283 NN (holotype, PMA; isotype, MO). Figure 2.

Although the species was validly published (Folsom, 1994a), a complete description of this species was not provided, and so one is offered here: Epiphytic, caespitose, roots 1-1.5 m diam.; stems to 30 cm long, 17-21 mm wide across conduplicate sheaths. Leaf blades $4.5-6.5 \times 0.7-1.4$ cm, oblong, obtuse, retuse or apiculate; flowers axillary, peduncles 15-20 mm long; floral bracts 11-14 × 8-14 mm. ovate-orbicular, acuminate, secondary bract lanceolate, ca. 11 × 2 mm; sepals and petals white or ivory with 7 to 9 lilac stripes, midlobe of lip lilac; sepals and petals ovate-lanceolate, narrowly acute-acuminate, margins hispid-ciliate; dorsal sepal 12-14 × 6 mm, lateral sepals $13-15 \times 6-7$ mm, petals 11-14 \times 4 mm; lip 10–12 \times 5–7 mm, claw 5.5–6 \times 3.5– 4 mm, dorsally keeled, midlobe 5 × 6 mm, lateral lobes 1-2.4 mm long; column 3.5-4 mm long, with distinct foot, with a rounded callus between column foot and lip, ligule 1.5-2 × 1.5-2 mm, subquadrate, shallowly emarginate; ovary 2-3 mm long, muricate.

Dichaea fragrantissima is readily distinguished from D. morrisii by the short, obtuse, lateral lobes of the ecallose lip, by the subquadrate ligule, and by the long, narrow, subacuminate sepals and petals. The subspecies fragrantissima is known only from cloud forests in Coclé and Veraguas provinces in Panama and has been collected only a few times. It differs from the new subspecies eburnea (described below) primarily in flower coloration and in having somewhat larger lateral lobes, or retrorse angles, on the lip. The specimens of subspecies fragrantissima we have seen have a prominent callus between the column foot and the claw of the lip, and the claw of the lip has a distinct median keel. Both of these features are variable within Panamanian collections of subsp. eburnea.

Additional specimens examined. PANAMA. Coclé: trail to La Mina, Hills N of El Valle de Antón, P. H. Allen 2874 (AMES), A. Maduro & E. Olmos 189 (MO, PMA).