



EPIDENDRUM ANDREI Hágster & L. Sánchez
THE GENUS EPIDENDRUM PART 2

Plate 306
ICONES ORCHIDACEARUM 3. 1999

EPIDENDRUM ANDREI Hágster et L. Sánchez, sp.nov.

Holotype: COLOMBIA: [Valle del Cauca]: Cauca, Tambores, Quindío, 15 Mar. 1976, E. F. André 2420 K! (illustration voucher).

Epidendro apagano Mansf. simile sed caulis anciptibus, sepalis obovatis, petalis spathulatis, labello obtrapezoidalii, clinandrio brevi eroso.

Hierba epífita, cespitosa, 14-29 cm de alto. Raíces basales, carnosas, delgadas, 1-1.5 mm de grosor. Tallos sencillos, tipo caña, anciptosos, ligeramente flexuosos, 8-24 x 0.5-1.5 cm. Hojas 4-6, equidistantes a lo largo del tallo, coriáceas; vainas anciptosas, lisas, 2-4 cm de largo; lámina elíptica, retusa a bilobada, carina dorsal baja, 5-10 x 2-3.5 cm. Espata ausente. Inflorescencia apical, en el tallo maduro, florece una sola vez, sésil. Brácteas florales mucho más cortas que el ovario, ovadas, agudas. Flores 2-3, resupinadas, simultáneas, verdes. Ovario ligeramente inflado ventralmente en la mitad apical, terete, 25-35 x 2-3.4 mm. Sépalos extendidos, glabros, libres, subcoriáceos, obovados, margen revoluto, 17-20 x 5-6 mm; el dorsal redondeado; los laterales oblicuos, agudos. Pétalos extendidos, espatulados, redondeados, margen extendido, membranáceos, 17-19 x 4-5 mm. Labelo unido a la columna, entero, base truncada, obtrapezoide al extenderse, margen entero, ápice entero, subcoriáceo, 10 x 12 mm; aparentemente ecalloso, carina baja, la cual se prolonga hasta el ápice. Columna recta, 9-10 mm de largo. Clinandrio corto, eroso.

Caespitose, epiphytic herb. Stems simple, reed-like, anciptous, slightly flexuose. Leaves 4-6, blade elliptic, apex retuse to bilobed. Inflorescence apical, sessile. Floral bracts much shorter than the ovary, ovate, acute. Flowers 2-3, resupinate, simultaneous, green. Ovary slightly inflated ventrally in the apical half, terete. Sepals spreading, glabrous, free, obovate, rounded, margin revolute; the laterals oblique, acute. Petals spreading, spatulate, rounded, membranaceous. Lip adnated to the column, entire, obtrapezoide; apparently ecallose. Column straight. Clinandrium short, eroso.

DISTRIBUTION AND ECOLOGY: Known only from the type from the Valley of the River Magdalena, epiphyte, in wet forest. Flowering in March.

RECOGNITION: *Epidendrum andrei* is member of the *E. difforme* Jacq. group and can be recognized by its anciptose stems, small, green flowers, sepals obovate, petals spathulate, lip obtrapezoid, short, erose clinandrium. It differs from *E. apaganum*, which has elliptic sepals and petals, 3-lobed lip, dentate lateral lobes, triangular, obtuse midlobe and a fimbriate-dentate clinandrium. *Epidendrum killipii* Hágster & L. Sánchez has larger plants, laterally compressed stems, larger flowers with elliptic sepals and rhombic petals, suborbicular-cordiform lip, subglobose callus and a prominent, dentate clinandrium.

CONSERVATION STATUS: DD (data deficient).

ETYMOLOGY: Dedicated to Edouard François André, who collected the material from which the type specimen was prepared, and presumably prepared the water color that is on the type sheet.

