



EPIDENDRUM BOYLEI Hágster & Dodson
THE GENUS EPIDENDRUM PART 3

Plate 420
ICONES ORCHIDACEARUM 4. 2001

EPIDENDRUM BOYLEI Hágster et Dodson, sp. nov.

Type: ECUADOR: Carchi: Cerro Golondrinas area, access via Chamorro property above El Carmen, which is above Hualchán. On peak of Cerro 2840. $00^{\circ} 50' N$ $78^{\circ} 12' W$, 2850-2860 m, 24 July 1993, **Brad Boyle & P. Hibbs 2307**. Holotype: MO! (illustration voucher).

Epidendrum manco Lindl. similare sed floribus respectu caulis erectis, labello trilobato, sepalis petalisque brunneis, labello albo-cremeo differt.

Hierba epífita, caespitosa, erecta, 8.5-10 cm de alto. Raíces basales, carnosas, delgadas, 0.5-1 mm de grosor. Tallos sencillos, tipo caña, rectos, anclíticos, 2.5-3.5 x 0.2-0.5 cm. Hojas 3-4 distribuidas en la mitad apical del tallo, erectas; vaina tubular, anclítica, estriada, 0.5-0.7 x 0.3-0.6 cm; lámina angostamente elíptica oblonga, desigualmente bilobada, con una quilla dorsal, margen apical crenado, coriácea, 3.0-3.5 x 0.6-0.8 cm. Espatas 1-2, tubulares, oblongas, anclíticas, subagudas, 1.8-2.0 x 0.4-0.5 cm. Inflorescencia apical, racemosa, erecta, 6 cm de largo; pedúnculo corto, terete a lateralmente aplanado, recto, no ornamentado. Brácteas florales mucho más cortas que el ovario o obsoletas. Ovario terete, no ornamentado, ligeramente dilatado hacia el ápice, 5-8 x 0.6-1.0 mm. Flores 4, sucesivas, no resupinadas, sépalos y pétalos pardos, labelo blanco cremoso; fragancia no registrada. Sépalos extendidos, ligeramente cóncavos, libres, obovados, agudos, margen entero, 5-nervados, 5.5-6.4 x 2.5-3.2 mm; el dorsal subredondeado. Pétalos extendidos, ligeramente cóncavos, libres, espatulados, obtusos, margen entero, 3-nervados, 4.8-5.5 x 1.5-2.0 mm. Labelo unido a la columna, reducido, trilobado, carnoso, 2.5 x 4.0 mm; bicalloso, con una quilla corta en medio, dando la apariencia de una estructura gruesa trilobada; lóbulos laterales diagonalmente subrómbicos, redondeados, margen finamente eroso; lóbulo medio semicircular, apiculado. Columna recta, 5 mm de largo. Clinandrio corto, entero. Antera no vista. Polinios no vistos. Rostelo subapical, hendido. Lóbulos laterales del estigma de 1/2 de largo de la cavidad estigmática. Nectario sin traspasar el perianto, no ornamentado. Cápsula no vista.

Epiphytic, caespitose herb. Stems cane-like, short, anclitose. Leaves 3-4, sheaths anclitose, blade oblong to narrowly elliptic, unequally bilobed, coriaceous, apical margin finely crenate. Spathes 1-2, tubular, narrow oblong, anclitose, subacute. Inflorescence apical, racemose, erect, few-flowered. Floral bracts much shorter than the ovary or obsolete. Ovary terete. Flowers 4, successive, non-resupinate, sepals and petals brown, lip creamy-white. Sepals spreading, free, obovate, rounded to acute, 5-veined. Petals spreading, spatulate, obtuse, 3-veined. Lip reduced, 3-lobed, fleshy, bicallose with a small intermediate short keel, producing a 3-lobed appearance; lateral lobes diagonally subrhombic, rounded, margin finely erose, midlobe semicircular, apiculate. Column straight. Clinandrium short, entire. Nectary short, not surpassing the perianth.

OTHER SPECIMENS: None seen.

RECOGNITION: *Epidendrum boylei* belongs to the mancum group and can be recognized by its small, caespitose plants with brown flowers and a creamy white lip, few flowers, anclitose stems, erect leaves, petals 3-veined, 3-lobed lip. It's similar to *E. mancum*, but the latter has orange and yellow flowers, elliptic-ovate, spreading leaves, entire lip. *Epidendrum brachyglossum* Lindl., is found at lower altitudes, plants are larger, have linear-lanceolate leaves, and a more or less 3-lobed lip.

DISTRIBUCION AND ECOLOGY: Known only from the type, epiphytic on shrub in páramo-elfín forest, 2850-2860 m. Flowering in July.

CONSERVATION STATUS: DD. Data deficient.

ETYMOLOGY: In honor of Brad Boyle, who collected the type specimen.

