



bracts infundibular, 3 mm long; pedicels 5 mm long; ovary trilamellate, 1.5 mm long and wide; *sepals* light orange, fleshy, concave, carinate, the dorsal sepal elliptical, obtuse, 11 mm long, 6 mm wide, 6-veined, free to the base, the lateral sepals oblong, oblique, connate 5 mm, 11 mm long, each 4.5 mm wide, 3-veined; *petals* oblong, obtuse, minutely apiculate, 4 mm long, 2.25 mm wide; *lip* thick, oblong, obtuse, 5 mm long, 2 mm wide, minutely spiculate with an obscure low, marginal lobe above the middle, with an apiculum beneath the tip, the disc with a pair of low, minutely serrated calli above the middle; *column* green, semiterete, 4 mm long, the apex bidentate, the foot stout, 3 mm long, deeply concave with a wide, circular margin, the pollinia a pair of irregular pollinia attached to the inner surface of the deciduous anther cap.

**Etymology:** Named for Bryon Rinke of Winfield, KS, who cultivates this species.

Among the various features that characterize this genus, *Kraenzlinella rinkei* is distinguished by a short ramicaul and a short raceme barely as long as the petiole of the leaf. The short, trilamellate ovaries are as long as they are wide. The fleshy, carinate flowers are nonresupinate. The short, oblong petals are minutely apiculate. The non-unguiculate lip is free of lobules, but a low pair of obscure, marginal lobes is present above the middle. Also above the middle the lip is minutely spiculate with a low pair of calli. A sharply acute apiculum is present beneath the tip.

***Kraenzlinella rinkei* Luer, sp. nov.** TYPE: COSTA RICA. Without collection data, flowered in cultivation in Winfield, Kansas, May 2010, by Bryon Rinke s.n. (Holotype: MO), C. Luer illustr. 21535. Fig. 19

*Planta mediocris, pedunculo racemo folio crasso petiolato multibreuiore, ovario brevissime lamellato, sepalis crassis carinatis, petalis late oblongis apiculatis, labello oblongo subtrilobato minutissime spiculato distinguitur.*

*Plant* medium in size, epiphytic, caespitose; roots coarse. *Ramicauls* relatively stout, erect to suberect, 2 cm long, enclosed by a tubular sheath equally long. *Leaf* erect, thickly coriaceous, elliptical-ovate, subacute, 9–10 cm long, 2 cm wide, cuneate below into a petiole 1–2 cm long. *Inflorescence* an erect, loose, successively few-flowered raceme up to 1.5 cm long including the peduncle 0.8 cm long, from the apex of the ramicaul at the base of the petiole, with a spathe 0.3 cm long; floral