



A new species of *Lecanorchis* (Orchidaceae) from Thailand

Somran Suddee⁽¹⁾ and Henrik Æ. Pedersen^(2*)

1. The Forest Herbarium (BKF), Department of National Parks, Wildlife and Plant Conservation, Chatuchak, Bangkok 10900, Thailand.

2. Botanical Garden & Museum, Natural History Museum of Denmark, University of Copenhagen, Gothersgade 130, DK-1123 Copenhagen K, Denmark.

* Corresponding author. Tel: +45 35 32 21 90; Email: henrikp@snm.ku.dk

(Manuscript received 9 August 2010; accepted 2 November 2010)

ABSTRACT: *Lecanorchis betongensis*, a new species from tropical rain forest in southern peninsular Thailand, is described and illustrated. The combination of semicircular column wings and a labellum with an odd number of major veins (but devoid of calli) places the new species in sect. *Lecanorchis*. However, the lack of any fusion between the labellum and the column readily distinguishes *L. betongensis* from all other species of the genus. The new species seems morphologically closest to *L. malaccensis* from Thailand, Malaysia and Sumatra.

KEY WORDS: Flora of Thailand, *Lecanorchis betongensis*, orchids, systematics, taxonomy, Vanilloideae.

INTRODUCTION

Due to its achlorophyllous condition, no cpDNA data are available for the mycoheterotrophic *Lecanorchis* Blume. However, phylogenetic studies based on morphological data (Freudenstein and Rasmussen, 1999) as well as sequence data from nuclear rDNA (Cameron and Chase, 2000) and mitochondrial DNA (Freudenstein and Chase, 2001) consistently indicate that this genus belongs to the Vanilloideae.

Morphologically, *Lecanorchis* is easily recognized from the presence of a cup-like calyculus just below the perianth, but the genus remains much understudied – mainly because it is only collected sporadically and because the fragile floral parts are often severely damaged when preserved by traditional pressing and drying.

According to the World Checklist of Selected Plant Families (<http://apps.kew.org/wcsp/home.do>; accessed on 1 September 2010), *Lecanorchis* accommodates ca. 18 species (probably an underestimate, as the leafless, dull-coloured plants constituting this genus are often overlooked in their natural environment). The generic distribution encompasses Japan (e.g. Hashimoto, 1990), Taiwan (e.g. Liu and Su, 1978), southern China (e.g. Chen et al., 2009), Vietnam (e.g. Averyanov, 2005), Bhutan and Sikkim (e.g. Pearce and Cribb, 2002), Thailand (see below), Peninsular Malaysia (e.g. Seidenfaden and Wood, 1992), Borneo (e.g. Wood and Cribb, 1994), Sumatra (e.g. Comber, 2001), Java (e.g. Comber, 1990), the Philippines (e.g. Valmayor, 1984) and New Guinea (e.g. Schlechter, 1914). The distribution map in Pridgeon et al. (2003: Fig. 221.2) additionally indicates

occurrence in Bangladesh, Myanmar, Laos, Cambodia, Sulawesi and Maluku, but we have been unable to find any explicitly validated records from these regions.

Seidenfaden (1978) recognized two species, *L. malaccensis* Ridl. and *L. multiflora* J.J.Sm., in his treatment of Thai *Lecanorchis*, and recently Suddee et al. (2010) reported *L. javanica* Blume and *L. nigricans* Honda as new records for Thailand. During the preparation of the Vanilloideae account for Flora of Thailand (Suddee, submitted), another species of *Lecanorchis* was discovered. The specimens collected from Betong District, Yala Province in peninsular Thailand near the Malaysian border, match no previously described species. Therefore, we propose the new species *L. betongensis*.

TAXONOMIC TREATMENT

Lecanorchis betongensis Suddee & H. A. Pedersen, *sp. nov.* Figs. 1 & 2

Type: Thailand, peninsular floristic region, Yala Province, Betong District, Ban Chulaphon Phattana 10, 500-600 m, 5 May 2007, *Wai 1053* (holotype PSU!, spirit).

Species haec L. malaccensis affinis, sed labello columna haud unito et sepalis petalisque multo minoribus differt.

Rhizome J-shaped. Roots fleshy, numerous, 2-12 cm long, 3-4 mm in diameter. Flowering shoots 20-65 cm tall, erect, dark purple to black, 3-4 mm in diameter at base, branched. Leaves reduced to sheaths, glabrous, ovate, 3-5 mm long, acute, coriaceous, amplexicaul at base. Inflorescence glabrous; rachis 1.5-4 cm long, 3- to 15-flowered; bracts ovate, glabrous, 0.5-1.5 by 1-1.5




Fig. 1. *Lecanorchis betongensis* Suddee & H. A. Pedersen. A: Flower. B: Column and labellum. C: Labellum. D: Petal. E: Dorsal sepal. F: Lateral sepal. G-H: Column. I: Fruits and flower bud. A-I: *Wai 1053*. Drawn by Pajaree Inthachup.

mm, margin membranous. Flowers long-pedicelled, green to light brown. Sepals glabrous; dorsal sepal oblong-ob lanceolate, 6-11 by 1.8-3 mm, 7-veined, apex hooded; lateral sepals oblong, 5.5-11 by 1.5-2.5 mm, 5-veined, apex hooded and apiculate. Petals glabrous, oblong, 5.5-10 by 2-3 mm, 3-veined. Labellum entire, rarely obscurely 3-lobed, 6-10 mm long along the midline, free from the column, 5-veined, major veins

branched above; side lobes not exceeding the column, indistinct; midlobe oblong-ob lanceolate, 2.5-3 mm wide, hooded, margin irregularly erose, with a small median hairy band, hairs multicellular. Column ventrally glabrous to finely puberulent, 4-8 mm long, winged at the top; wings semicircular with finely papillose edges; anther ca. 1 mm long, muricate at apex. Ovary (including pedicel) 8-20 mm long; calyculus




Fig. 2. *Lecanorchis betongensis* Suddee & H. A. Pedersen. A: Underground organs. B: Habit. C: Flower. D: Flower (sepals and petals removed). E: Labellum. F: Column. A–F: *Wai* 2003. Photographed by Thamarat Phutthai (A) and Jarearnsak Sae Wai (B–F).

glabrous, ca. 0.5 mm long, margin irregularly denticulate. *Capsule* black, fusiform, 2–2.3 cm long, 2–2.2 mm in diameter.

Specimens of *L. betongensis* examined (all from Thailand, Yala Province, Betong District): Ban Chulaphon Phattana 10, 500–600 m, 15 August 2004, *Wai* 69 (PSU, spirit); 5 May 2007, *Wai* 1053 (holotype PSU, spirit); 17 September 2010, *Wai* 2003 (PSU, spirit); Nikhom Betong-Aiyoe Khwin, 700 m, 26 February

2001, *Suksathan* 2903 (QBG).

Specimens of *L. malaccensis* examined for comparison: Peninsular Malaysia: Pahang, Tahan Woods, 1891, *Ridley s.n.* (SING, type). – Thailand: northeastern floristic region: Loei Province, Phu Kradueng, trail to Pha Lom Sak, 1220 m, 15 March 2008, *Suddee, Puthai, Kerdkaew, Witsawachaiwat, Hemrat, Somngam & Suwanachat* 3584 (BKF, spirit); Phu Luang, Lon Tae, 1215 m, 11 November 2008, *Suddee, Pedersen, Watthana, Puthai, Hemrat & Suwanachat* 3988 (BKF, spirit). Peninsular floristic region:



Narathiwat Province, Sukhirin, 29 March 2000, *Puudjaa & Cholkulchana* 737 (BKF, spirit); Surat Thani Province, Ko Pha-ngan, 2 June 1927, *Pit* 761 (C, spirit).

Etymology: The new species is named after the Betong District.

Distribution: *Lecanorchis betongensis* is apparently endemic to Thailand, but it might be expected also to occur across the border in Peninsular Malaysia.

Conservation: Known only from 4 collections from the type locality and a nearby locality within the same subdistrict. The area is in Bang Lang National Park; it is well protected, but its flora is still poorly known.

Ecology: Mycoheterotrophic, apparently restricted to tropical rain forest at 500-700 m alt. Flowering: February-September.

Notes: The new species seems morphologically closest to *L. malaccensis* from Thailand, Malaysia and Sumatra. However, it differs from the latter species in having larger sepals and petals, and in its labellum being completely free from the column. Indeed, the lack of any fusion between the labellum and the column readily distinguishes *L. betongensis* from all other species of the genus. The combination of semicircular column wings and a labellum with an odd number of major veins (but devoid of calli) places the new species in sect. *Lecanorchis* as morphologically defined by Hashimoto (1990). However, the entire or only obscurely 3-lobed condition of the labellum in *L. betongensis* is not typical of sect. *Lecanorchis*.

ACKNOWLEDGEMENTS

Thanks are due to Jarearnsak Sae Wai for photographs, for useful information and for discussions about localities, to Pajaree Inthachup for preparing the line drawing and to Kitichate Sridith and PSU staff for making collections and facilities available during the first author's visits.

LITERATURE CITED

- Averyanov, L. V.** 2005. New orchids from Vietnam. *Rheedea* **15**: 83-101.
- Cameron, K. M. and M. W. Chase** 2000. Nuclear 18S rDNA sequences of Orchidaceae confirm the subfamilial status and circumscription of Vanilloideae. In: Wilson, K. L. and D. A. Morrison (eds.), *Monocots: systematics and evolution*: 457-464. CSIRO, Melbourne, Australia.
- Chen, X., P. J. Cribb and S. W. Gale** 2009. 55. *Lecanorchis* Blume, Mus. Bot. 2: 188. 1856. In: Wu, Z. et al. (eds.), *Flora of China* **25**: Orchidaceae: 171-172. Science Press, Beijing & Missouri Botanical Garden, St. Louis, USA.
- Comber, J. B.** 1990. *Orchids of Java*. Bentham-Moxon Trust. Royal Botanic Gardens, Kew, UK. 407pp.
- Comber, J. B.** 2001. *Orchids of Sumatra*. Royal Botanic Gardens, Kew, UK. 1026pp.
- Freudenstein, J. V. and M. W. Chase** 2001. Analysis of mitochondrial *nad1b-c* intron sequences in Orchidaceae: utility and coding of length-change characters. *Syst. Bot.* **26**: 643-657.
- Freudenstein, J. V. and F. N. Rasmussen** 1999. What does morphology tell us about orchid relationships? – A cladistic analysis. *Am. J. Bot.* **86**: 225-248.
- Hashimoto, T.** 1990. A taxonomic review of the Japanese *Lecanorchis* (Orchidaceae). *Ann. Tsukuba Bot. Gard.* **9**: 1-40.
- Liu, T.-S. and H.-J. Su** 1978. 38. Orchidaceae. In: H.-L. Li et al. (eds), *Flora of Taiwan* **V**: 859-1137. Epoch Publishing Co., Taipei, Taiwan.
- Pearce, N. R. and P. J. Cribb** 2002. *Flora of Bhutan including a record of plants from Sikkim and Darjeeling* 3(3). The orchids of Bhutan. Royal Botanic Garden, Edinburgh, UK & Royal Government of Bhutan, Thimbu, Bhutan. 643pp.
- Pridgeon, A. M., P. J. Cribb, M. W. Chase and F. N. Rasmussen** 2003. *Genera Orchidacearum* 3. Orchidoideae (part 2), Vanilloideae. Oxford University Press, Oxford, UK. 358pp.
- Schlechter, R.** 1911-1914. *Die Orchidaceen von Deutsch-Neu-Guinea*. *Beih. Repert. Spec. Nov. Regni Veg.* **I**: 1-1079.
- Seidenfaden, G.** 1978. Orchid genera in Thailand VI. Neottioideae Lindl. *Dansk Bot. Arkiv* **32(2)**: 1-195.
- Seidenfaden, G. and J. J. Wood** 1992. *The orchids of Peninsular Malaysia and Singapore. A revision of R. E. Holttum: Orchids of Malaya*. Olsen & Olsen, Fredensborg, Denmark. 779pp.
- Suddee, S., S. Chantanaorrapint, P. Tripetch and S. Thainukul** 2010. New records in *Lecanorchis* Blume and *Vanilla* Plum. ex Mill. from Thailand, with keys to the Thai species. *Thai Forest Bull. (Bot.)* **38**: 1-7.
- Valmayor, H. L.** 1984. *Orchidiana Philippiniana*. Eugenio Lopez Foundation, Manila, Philippines. 737pp.
- Wood, J. J. and P. J. Cribb** 1994. A checklist of the orchids of Borneo. Royal Botanic Gardens, Kew, UK. 409pp.



泰國的新種皿蘭屬(蘭科)植物

Somran Suddee⁽¹⁾ and Henrik Æ. Pedersen^(2*)

1. The Forest Herbarium (BKF), Department of National Parks, Wildlife and Plant Conservation, Chatuchak, Bangkok 10900, Thailand.

2. Botanical Garden & Museum, Natural History Museum of Denmark, University of Copenhagen, Gothersgade 130, DK-1123 Copenhagen K, Denmark.

* 通信作者。Tel: +45 35 32 21 90; Email: henrikp@snm.ku.dk

(收稿日期：2010年8月9日；接受日期：2010年11月2日)

摘要：本文敘述並描繪泰國南部半島熱帶雨林的新種 *Lecanorchis betongensis*。該新種具有半圓形的蕊柱側翼，且唇瓣具奇數主脈（但缺乏肉突），故應屬於 *Lecanorchis* 節。然而，其唇瓣與蕊柱完全分離而使它與同屬其他物種明顯不同。這個新種形態上與泰國、馬來西亞、蘇門答臘的 *L. malaccensis* 相近。

關鍵詞：泰國植物誌、*Lecanorchis betongensis*、蘭花、系統學、分類學、香莢蘭亞科。