

Lepanthes gerardensis M.A. Blanco, *sp. nova*

TYPE: COSTA RICA. Puntarenas: Monteverde, Valle de San Gerardo, entre Santa Elena y Arenal, vertiente atlántica, ca. 10°23'00"N, 84°47'50"W, 1000 m, 23 enero 1999, M. Blanco & V. Arias 924 (holotype, USJ).

FIG. 1

Lepanthes ancylopetalae Dressler primo aspectu similis, sed habito minore, foliis petiolatis, late ellipticis, labello relative magno, bilaminato, cum laminis glabris, deltoideis, una contra alteram dispositis difert.

Plant a caespitose epiphytic herb. *Roots* up to 5 cm long, 0.5 mm thick, terete. *Ramicauls* erect, terete, 1-10 mm long, 0.3 mm thick, clasped by 1-4 cataphylls, with a single terminal leaf. *Cataphylls* 1-4 mm long,

pale brown (dry at maturity), lepanthiform, glabrous to minutely ridged, acute apically; ostia oblique and expanded (up to 1 mm in diameter). *Leaves* petiolate, thick, green with purple dots on the abaxial surface. *Petiole* 1-3 mm long, 0.3 mm thick. *Blade* conduplicate, widely elliptic when spread, 7-9 mm long, 5-6 mm wide, about 1 mm thick, cuneate basally, obtuse apically, mucronate, with 7 longitudinal veins, margin entire. *Inflorescence* terminal on the ramicaul, a pedunculate raceme; peduncle terete, 4 mm long, 0.1 mm thick; the raceme up to 3 mm long, distichous, sequentially flowered (up to at least 15 flowers produced). *Flower bracts* triangular, 0.5 mm long. *Flowers* showy, large for the plant, resupinate; sepals pale greenish yellow, petals yellow with upper lobe and inner margin of lower lobe red, lip and column dark red, anther and appendix white. *Dorsal sepal* ovate-lanceolate, acute, 3-veined, 3.0 mm long, 1.5 mm wide, glabrous, margin entire. *Lateral sepals* obliquely lanceolate, 2-veined, connate for half their length, acute, each 2.5 mm long and 1.5 mm wide, glabrous, margin entire. *Petals* two-lobed, with a minute median apicule, minutely papillose, 1.5 mm wide, 0.6 mm long; the upper lobe oblong, rounded apically, the base expanded, 1 mm long; lower lobe subtriangular, rounded, 0.4 mm long; glabrous, margin entire. *Lip* bilaminate, surrounding the column;

blades rounded-deltoid, with the angles directed inwards, minutely papillate, each 1 mm long, 0.5 mm wide, margin entire; appendix digitiform, pointing toward stigma, minutely ciliate, 0.2 mm long. *Column* clavate, slightly curving downward, fused with the lip at its base, 1 mm long, anther terminal, 0.2 mm long; stigma ventral, partially bilobed. *Pollinarium* not seen. *Fruit* not seen.

Lepanthes gerardensis can be recognized by its short habit (less than 2 cm tall), widely elliptic leaves, relatively large flowers, and petals with a small lower lobe and red upper lobe. Otherwise, the flower morphology is typical of most species in the genus.

Its flowers are most similar to those of the Mexican *L. ancylopetala* Dressler, but that species has a concave dorsal sepal and a smaller lip with pilose blades; furthermore, plants of *L. ancylopetala* are taller (up to 6 cm tall) with narrowly elliptic leaves (figure 8 in Salazar-Chávez & Soto-Arenas 1996). Vegetatively, *L. gerardensis* resembles *L. papillipetala* Dressler, also from Mexico (figure 54 in Salazar-Chávez & Soto-Arenas 1996).

Behar & Tinschert (1998: 96) present a color photograph of a Guatemalan *Lepanthes* identified as *L. papillipetala* (the same flower and the whole plant are also illustrated in Balistreri 1998: 53); this is consistent with Luer's interpretation of *L. papillipetala* (C. Luer, pers. commun. 2003). However, it is different from the interpretation of *L. papillipetala* in Salazar-Chávez & Soto-Arenas's (1996) revision of the Mexican species of *Lepanthes*; it is more similar to their illustration of *L. ancylopetala*, and even more similar to *L. gerardensis*, at least in floral shape and color. Because Salazar and Soto not only studied the types of both species, but also studied populations of the type locality in Chiapas (the same for both *L. ancylopetala* and *L. papillipetala*), I choose to follow their taxonomic interpretations.

Among its Costa Rican congeners, *Lepanthes gerardensis* resembles *L. excedens* Ames & Correll (= *L. empis* Luer) in plant size and leaf shape. The flowers are superficially similar, but the petals of the latter species are crescentiform, with angular upper and lower lobes, and the lip has narrow (non-peltate) strongly arching blades continuous with the connectives, as well as a stouter appendix.

ETYMOLOGY. Named after the San Gerardo Valley near Monteverde, where the species was found.

ECOLOGY: Apparently endemic to the Caribbean slopes of the Tilarán Cordillera of Costa Rica, in cloud forest at around 1000 m elevation.

Many plants were found growing on guava trees (*Psidium guajava*, Myrtaceae) in a pasture adjacent to primary forest. It appears to be a specialized twig epiphyte; the many plants observed were restricted to terminal branches of 1 cm or less in diameter, almost devoid of mosses or liverworts. Other well-known twig epiphytes such as *Erycina* (*Psylgmorchis*) *pusilla*, *Epidendrum vincentinum*, and *Hybochilus inconspicuus* also occurred on the same twigs.

Flowering occurs at least in January in the field, and possibly throughout the year. Several plants brought to Lankester Botanical Garden survived for about two years, flowering constantly under more or less uniform humidity conditions.

ACKNOWLEDGMENTS: I am grateful to Jorge Gómez-Laurito (USJ) for the Latin diagnosis and Vernon Arias for companionship in the field. Suggestions by Carlyle Luer and Alec Pridgeon greatly improved the manuscript.




Figure 1. *Lepanthes gerardensis* M.A. Blanco. A – Habit. B – Flower. C – Dissected perianth. D – Lip and column, lateral view. ILLUSTRATION VOUCHER: Blanco & Arias 924 (USJ).