

domonopodium formantes. Folia disticha. Inflorescentiae uniflorae. Sepala calcarata. Labellum sepalorum calcari intus adnatum. Columna apoda. Rostellum prominens, aciculare. Stipes polliniorum abbreviata.

Genera: *Cryptocentrum* Bentham.

Anthosiphon Schlechter.

Notylia Norae *Garay* sp. nov.

Epiphytica, pusilla, ca. 2.5 cm. alta; radicibus filiformibus, flexuosis, glabris; foliis equitantibus, oblique ovato-lanceolatis, carnosis, 5–8 mm. longis, 1.5–2 mm. latis; inflorescentia erecta vel paulo arcuata, pauciflora; bracteis ovatis, acutis, ca. 1 mm. longis; floribus diaphanis, lilacinis; sepalo postico ovato, conduplicato, apice acuto, 1-nervio, 3 mm. longo, 1.7 mm. lato; sepalis lateralibus liberis, anguste oblongo-lanceolatis, acutis, 1-nerviis, 4 mm. longis, 0.8 mm. latis; petalis ovato-lanceolatis, acutis, 1-nerviis, 4 mm. longis, 1 mm. latis; labello longe unguiculato, deinde in laminam ovatam in medio leviter constrictam expanso, apice acuto, basi minutissime ciliato, 3-nervio, 3.5 mm. longo, antice 1.8 mm. lato; columna erecta, capitata, ca. 2 mm. longa; ovario cum pedicello ca. 2.5 mm. longo.

Notylia Norae is related to the Bolivian *N. lilacina* Kränzl. but differs from it in the shape of the petals and lip.

It gives me great pleasure to dedicate this beautiful miniature orchid to Mrs. Dunsterville, whose inspiration and special interest in botanicals has been of indispensable help in the preparation of the Iconography of the Venezuelan Orchids.

VENEZUELA: Kavanupa, Rio Uaiparu. On small tree near water's edge. G. C. K. Dunsterville 442 (TYPE in Herb. Ames).

Telipogon Teuscheri *Garay* sp. nov.

(§ *Brevicaules*). Radicibus satis longis, copiosis; cauli-