MASDEVALLIA ABBREVIATA, n. sp.* Another member of the group of the Amandæ. I have had a day's work to go once more over all my

types of the affinity of Masdevallia polysticta, melanopus, x anthodactyla, Lehmanni, tridens, caloptera, Gustavi, pachyura. This one is the ninth species, so much a connecting link between M. polysticta and melanopus that I often believed it to be a mule between these two. It has the full colours of M. polysticta, yet there are usually far fewer dots. There is even a nearly unspotted variety. The whole flower is far shorter, more gibbous underneath at its base,

and those cilie and hairs, so very conspicuous in Masdevallia polysticta, are scarcely to be seen, or not at all present. Then the petals are more retuse, with

a single apiculus, and very seldom accrned with a few lateral teeth (only seen once). The anterior part of the lip is hooded, as in genuine Masdevallia polysticta, and covered with three blunt ridges, yet it is usually far broader. I am not very sure whether it comes from M. Roezl, or perhaps from M. Bruchmüller, since none of wild specimens agree with it. I am rather more inclined to believe in its Roezlian origin. Masdevallia Amanda has the same long lateral sepals, which are charateristic of M. polysticta, and which I saw very judiciously represented in an elaborate drawing of this species prepared for the Botanical Magazine by Mrs. Dyer, H. G. Rekb. f.

^{*} Masdevallia abbreviata, n. sp.—(Amandæ.) Folius spatulatis obtusis; racemo paucifloro; bracteis cucullatis ovaria crispo-altat vulgo superaotibus; perigonio abbreviato ante mentum obtusum constricto; sepalorum partibus latis brevismisti trianquilis minutissime denticulatis, in caudas longiores simisti transpulsi minutissime denticulatis, in caudas longiores apriculto medio, labelli lachnis lateralibus semioblogis cloragatis, lamma antica infraposicia oblonga lateribus replicati, lineis tumidis ternis: columna gracileota, androclinii cucullo serrulato. H. G. Rehb. J.