

MASDEVALLIA Ruiz & Pav.

M. (§ Triangulares) **TUBULIFLORA** sp. nov. Aff. *M. floribundæ* Lindl. *Caules* secundarii erecti vel erecto-patuli, 1 cm. longi. *Folium* oblanceolatum, obtusum, rigidum, 5–8 cm. longum, 9–13 mm. latum. *Pedunculus* gracilis, filiformis, folio brevior. *Flores* purpureo-striati?, ± 1 cm. longi. *Sepalis* usque ad medium coalitis. *Calyx* tubulosus; lobi laterales falcato-lanceolati, 6 mm. longi, 3-nervii; lobus dorsalis carnulosus, linearis, obtusus, 7 mm. longus. *Petala* lineari-lanceolata, uninervia, 4 mm. longa, obtusa, apiculata. *Labellum* lineare, subacutum, ad basim cordatum, lateraliter ad medium bicallosum, 4 mm. longum, 1 mm. latum. *Columna* 3.5 mm. longa.

The petals are dilated at about the middle and provided with a blunt lobule on the anterior margin, 2 mm. from the base. The flowers appear to have been whitish or yellowish when fresh and striped with purple in the throat of the calyx.

Alt. 350 m., Cubilquitz, Depart. Alta Verapaz, Guatemala, January, 1902, *H. von Tuerckheim* (no. 512).