

2. *N. LINDLEYI*, n. sp.—Folia anguste lanceolata, acuminata, petiolata. Scapus bipedalis v. altior. Racemus elongatus, puberulis, multiflorus. Bracteæ anguste lanceolatæ, puberulæ. Ovarium puberulum, triquetro-oblongum. Sepala lineari-lanceolata, minute puberula, cum petalis et labellum cuspidatis. Petala sublatis, extus carinata. Labellum petalis subsimile, callo medio lineare paullo incrassato. Filamenta ultra dimidium libera. Capsula ovoideo-oblonga, triquetra, subglabra.

Hab. Borneo; *Lowe!* Island of Penang; *Curtis*, n. 469!

A tall plant, reaching to 3 or $3\frac{1}{2}$ feet high. Leaves 1–2 feet long by $1\frac{1}{2}$ –2 in. broad. Racemes 10–15 in. long, with numerous flowers. Bracts $\frac{3}{4}$ – $2\frac{1}{2}$ in. long by $1\frac{1}{2}$ –3 lin. broad. Pedicels 1–2 lin. long. Ovary 4–5 lin. long. Segments 7–9 lin. long. Column 1 lin., free portion of filaments $1\frac{1}{2}$ lin. long, anthers $3\frac{1}{2}$ lin. long. Free portion of style 3 lin. long. Capsule $\frac{1}{2}$ in. long.

Evidently allied to the preceding, but with narrower leaves, less pubescence, and other differences. Lindley appears to have considered it identical with *N. Zollingeri*, but, as I think, quite wrongly. Likewise Reichenbach, who (*Bonpl.* v. p. 58) remarks:—"In herbario Lindleyano adest planta quæ omnino *N. Zollingeri* bene evoluta. Inflorescentia prope *Cœlicæ macrostachyæ* seu *Calanthidis* cujusdam. Ovarium abrupte turbinatum; apice recurrens in rostrum. Sepala linearia, elongata (ovario incluso callo longiora) apicibus apiculata, apiculis in carinulis exeuntibus. Tepala subbreviora. Labellum prope ejusdem rationis, convexum, pagina inferiori carinatum.—Sepala oblique inserta. Stylus apice retusus; filamenta lateralia extus decurrentia. Antheræ lineares apice obtuse acutæ basi cordatæ; versatiles. Borneo, *Lowe.*" But the sheet referred to in Lindley's Herbarium contains a single specimen of *N. Zollingeri*, Reichb. f. (collected by Zollinger himself, in Java), and an enlarged drawing of a single flower, labelled by Lindley himself—"Borneo, *Lowe*, in Hb. Hooker." This drawing, so fully described by Reichenbach, is from the very specimen now described by me as *N. Lindleyi*, a quite distinct plant from *N. Zollingeri*, Reichb. f., as remarks under that species will show. The Penang plant seems identical in every respect with the Bornean one; and being in much more perfect condition, I have made use of it in drawing up the description wherever the other was insufficient.