

2. ZYGOPETALUM CERINUM. Huc: *Huntleya cerina* Lindl. Paxt. Fl. G. III. (1851). 62:

Sepalis subrotundis concavis, labello ovato convexo, retuso, crista
crassa semicirculari truncata plicata, columna apice nuda.

82*

Firm, whitish, waxy blossoms, not unlike those of *Maxill. Harrisoniae*, *Veraguas* by Warscewicz. — Chiriqui Volc. 8000'. General appearance of the *Huntleya violacea*. Flowers rise singly from the base of the leaves upon a peduncle about 6' long with a few short tubular bracts 3 inches across, the sepals are rounded, and even at the edge, of a very pale straw colour. Lip is somewhat ovate, convex, indented at the point, much more yellow and furnished near the base with a deep thick semicircular ruff, composed of numerous plaited folds. The column is deep violet near the base, has no expansion or hood over the anther.

Huc: *Pescatoria* Rchb. fil. in v. Schl. Bot. Ztg. 1852. p. 667:

Perigonii subcarnosi phylla 5 cuneato-obovata, acutiuscula, tria superiora recta, lateralia externa oblique in pede labelli inserta. Labelum unguiculatum, unguis ligulatus, media basi carinatus carina triangula, basi utrinque triangule auriculatus, sub angulo obtuso in basin gynostemii excavatam introrsus (uti ipsissima labelli basis in *Miltonia spectabili*). Lamina triloba, lobi laterales trianguli, erecti, disco carnoso utrumque conjungente; discus medio longitudinaliter carinatus, antrorum rotundatus, multicristatus, cristulae obtusatae, oblique imbricatae; lobus labelli medius rotundatus revolutus, ipso limbo revoluto a lobis lateralibus erectis separatus. Gynostemium aequaliter semiteres, antrorum excavatum; androclinium prope perpendiculare, apice productum, immarginatum; processus rostellaris trilobus, dens medius linearis, corneus, lobi laterales membranacei, integri, stigma tegentes. Stigma ellipticum, limbo inferiore trilobulato. Apparatus pollinis Zygo-petali.

Genus dicatum prae nobilissimo Pescatore, Orchidearum cultori celeberrimo. **P.**
cerina (*Huntleya cerina* Lindl.)