

215. *Tillandsia alata*, Kuhn, ad *T. fusca*, Lindl.,
accedit, recedit lacinias microribus, petalis et lobis
laciniosis.

Pseudobulbi suborbiculari-oblongi, 2-3 lin. longi. *Folia*
lineari-oblonga, acuta, canaliculata, valde varinata, $2\frac{1}{2}$ poli.
longa, 1- $1\frac{1}{2}$ lin. lata. *Pseudobulbi* breviter, unilobati. *Fructus*
lanco-lanceolatus, acutus, 3 lin. longus. *Pedicelli* obliquiter-alati,
10-12 lin. longi. *Sepala* lanceolato-acuta, acuta, subvarinata,
7 lin. longa, $2\frac{1}{2}$ lin. lata. *Petalis* obovato-oblonga, obtusa, $6\frac{1}{2}$ lin.
longa, $1\frac{1}{2}$ lin. lata. *Lobellae* orbiculari-obovatum, retusum,
longe reflexissimum, 8 lin. longum, $6\frac{1}{2}$ lin. latum. *Calcar* 10 lin.
longum, attenuatum. *Columnae* lata, $1\frac{1}{2}$ lin. longa, alba lacin
oblongis subintegra.

COLOMBIA : Millium.

Flowered in the collection of Sir Trevor Lawrence, Bart., in
June, 1835. Flowers white, except the basal two-thirds of the
lip, which is deep yellow with seven light red veins along
the centre, and the spur, which is yellowish.