CCXLVIII.—NEW ORCHIDS: DECADE 2.

11. Pleurothallis subulata, Rolfe, n. sp.—Caules graciles. Folia cordato-oblonga, apice attenuata, subobtusa, $4\frac{1}{2}$ poll. longa, $1\frac{1}{2}$ poll. lata. Flores solitarii. Spatha lanceolata, 6 lin. longa. Pedicellus 6-7 lin. longus. Sepalum posticum ovato-oblongum, acutum, 6 lin. longum, 4 lin. latum. Sepala lateralia omnino connata, ovato-oblonga, subacuta, postico æqualia. Petala subulata, 3 lin. longa. Labellum subpandurato-ovatum, acutum, 3 lin. longum, $1\frac{1}{4}$ lin. latum. Columna brevis.

138

A species belonging to Lindley's section Macrophyllæ fasciculatæ, which flowered in the collection of Baron Hruby, of Peckau, Bohemia, and was sent to Kew for determination through Messrs. F. Sander & Co., of St. Albans. Lindley separated the species with flowers over half an inch long from those with smaller flowers, under the name Grandi-floræ, and into this group the present species would fall. But I believe this arrangement is purely artificial, and were it abolished, P. subulata would be placed near to P. cardiothallis, Rchb. f., which it closely resembles in several respects, though its flowers are far larger. Like several of its allies, it has flowers of a dark lurid purple. Its native country is not recorded. The name refers to the petals.