

5. ***PLEUROTHALLIS GIRALDOI*** Luer, sp. nov.


Planta grandis, folio plano profunde cordato, spatha grandi suberecta, sepalo dorsali synsepaloque late ovatis apice rotundatis, petalis oblongis erosis, labello ovoideo apice rotundo convexo marginibus erosis distinguitur.

COLOMBIA: Dept. of Valle del Cauca: Finca Zíngara, between Cali and Buenaventura, alt. 1900 m, 15 Jan. 1994, J. Giraldo G. & L. Olver A. 97 (Holotype: TULV), C. Luer illustr. 18414; near Bitaco, km. 18 on the old road from Cali to Buenaventura, alt. 1800 m, collected by J. L. Aguirre, 1993, cultivated at Colomborquídeas, flowered in cult. 21 October 1997, R. Escobar 8301 (JAUM).

ETIMOLOGÍA: Dedicada a Jorge Giraldo G., de la Universidad del Valle en Cali, Colombia, co-descubridor de la especie.

ETYMOLOGY: Named in honor of Jorge Giraldo G. of the Universidad del Valle, Cali, Colombia, co-discoverer of this species.

Plant large, epiphytic, caespitose; roots slender. Ramicauls slender, erect, 17-27 cm long, with a loose, tubular sheath on the lower third and 2 other sheaths at the base. **Leaf** coriaceous, spreading, cordate-ovate, obtuse, shortly acuminate, 9-11 cm long, 5-7 cm wide, the base sessile, deeply cordate with the lobes 2 cm deep. **Inflorescence** a fascicle of successive, single, red-purple flowers, borne from a suberect, conduplicate spathe at the base of the leaf, 14-18 mm long, 5-6 mm broad, enclosing the peduncles, pedicels and floral bracts; peduncles 2-3 mm long; floral bracts tubular, 7 mm long; pedicels ca. 10 mm long; ovary 5 mm long; **sepals** cellular-glandular within, glabrous externally, the dorsal sepal broadly ovate to suborbicular, rounded at the apex, 11 mm long, 10 mm wide, 3-(5)-veined, the lateral sepals connate into an ovate synsepal with the apex rounded, 11 mm long, 9.5 mm wide, 4-(6)-veined; **petals** oblong, denticulate-erose, cellular-glandular, acute, oblique at the base, 8 mm long, 1.5 mm wide, 1-veined; **lip** purple, glabrous, ovoid, convex, with the sides of the basal thirds recurved, 5 mm long, 5.25 mm wide, with the margins denticulate-erose, with the apex rounded with a tubercle beneath, with the surface of the convex apical half indistinctly demarcated into a low, rounded prominence, the base deflexed into a flat claw in apposition to the column-foot, hinged



5. *PLEUROTHALLIS GIRALDOI* Luer.

to the base of the column-foot; **column** stout, 1.5 mm long, 2.75 mm broad, the foot thick, obsolescent, with the anther, rostellum and stigma apical.

Esta especie, del subgénero *Pleurothallis*, sección *Macrophyllae-Fasciculatae*, pertenece a la subsección *Cardiostolae*, por la espata notable, suberecta. La hoja es grande, patente, profundamente cordata. Las flores son bilabiadas; los sépalos redondeados en el ápice, y de escasa venación para su tamaño; los pétalos oblongos, erosos; el labelo convexo, con los bordes revolutos en su tercera parte basal.

This species of subgenus *Pleurothallis* section *Macrophyllae-Fasciculatae* and subsection *Cardiostolae* by reason of the conspicuous, suberect spathe. The leaf is large, deeply cordate and spreading. The sepals of the bilabiate flower are sparsely veined for their size, rounded at their apices; the petals are oblong and erose; and the lip is convex with the sides of the basal thirds revolute.