

white without, with green apices. *Petals* white or cream with a green median vein without; midlobe of lip yellow. *Lip* with a purplish submarginal band on each side; base of lip minutely striped with white and brown within; sepaline tube 4-8 mm. *Dorsal sepal* elliptic-oblong to oblanceolate, obtuse, apiculate, 3.5-5 x 1.1-1.5 cm. *Lateral sepals* ovate-lanceolate or oblanceolate, obtuse, apiculate. *Petals* narrowly obovate or oblong-elliptic, obtuse, 2.8-4.7 x 1-1.5 cm. *Lip* subquadrate-subobovate, 3.2-4 x 3-3.8 cm, basal calli 7-9 mm; base of lip with ca. 23 low ridges, these becoming higher and more ornate distally. *Column* 20-27 cm x 4.5-6 mm laterally, 5-5.5 cm. dorsoventrally, lateral arms 1-2 mm.

This species is one of the most distinctive known to us. It does not resemble any other species known from Central America. It is, in fact, one of the most difficult to deal with. The base of the lip has about 22 low keels. The keels and the alternating grooves appear as narrow stripes. The keels become much higher in the distal portion of the lip, but they are actually soft and mealy. It is difficult to handle the lip without rubbing off the “keels.” All *Sobralia* flowers are delicate, but it is almost impossible to handle the lips of this species without destroying the surface. Thus, we have chosen the epithet “*fragilis*” for this species. It is quite possible that this species is related to a Colombian species that has been called “*S. suaveolens*.” The so-called “*S. suaveolens*” has prominent “crests” on the lip that are actually a powdery, yellow substance, and may act as pseudopollen to attract female bees that gather the pseudopollen as food for their brood. This Colombian species has the lip very much more deeply divided than either *Sobralia fragilis* or the true *S. suaveolens*. It is quite possible that *S. fragilis* and the misidentified Colombian species are related to each other and they may both attract pollinators by pseudopollen.

We have only two plants of *S. fragilis*, but the species clearly show synchronized flowering, with both plants usually flowering on the same days.

DISTRIBUTION: only known from Costa Rica.

ETYMOLOGY: from the Latin *fragilis*, “fragile”, referring to the delicate, easily damaged surface of the lip.

Sobralia geminata* Dressler & Bogarín, *sp. nov.

TYPE: COSTA RICA. Cartago: Paraíso, Cachi, Peñas Blancas, camino a Cerros Duán, 9°49'51.3"N 83°46'13.1"W, 1400 m, bosque muy húmedo premontano, terrestre cerca del camino, flowered in cult. 1 October 2009, R.L. Dressler 7173 (holotype: CR; isotype: JBL-spirit). FIG. 5, 6, 8C.

A *Sobralia chrysostoma* Dressler similis, floribus minoribus, labello maculis brunneis plerumque aliquantum albo notato recedit.

Roots 5-7 mm in diameter; stems 20-140 cm, basally 5-6 mm in diameter, distally 3.5-5 mm in diameter. *Stems* dark greenish purple mottled with pale green, young sheaths similar, especially opposite the next leaf blade. *Leaves* elliptic or broadly elliptic, acuminate, apiculate, 11-25 x 4-7.4 cm, with 7-9 raised veins beneath. *Inflorescence* ellipsoid, 3.5-4 x 0.7-1 cm; smaller foliar bracts 3-10 x 1.5-5 cm; ovary 10-22 mm, sepaline tube 1-2 cm. *Sepals* greenish cream with green mid-veins, apically green. *Petals* white without, lip yellow with white margins and usually some white on disk, disk with pale brown spots or streaks. *Dorsal sepal* oblong-elliptic or oblanceolate-elliptic, apiculate, 4.2-6.6 x 1.25-1.8 cm. *Lateral sepals* oblong-elliptic or narrowly obovate, acute, 4.5-6 x 1.5-2.3 cm. *Petals* oblong-elliptic or narrowly obovate, subobtusate to subacute; 4.3-7.1 x 1.6-2.5 cm. *Lip* obovate, 5.3-6.5 x 4.2-4.8 cm, basal calli 0.8-1.8 cm, median keels 4-9, low, sometimes with 1-2 shallow grooves, midlobe crisped. *Column* 2.8-4 cm x 5-9 mm laterally x 5-8 mm dorsoventrally; lateral lobes (arms) small, tooth-like, 1.5-1.6 mm.

PARATYPES: Costa Rica. **Alajuela:** San Ramón, Piedades, Piedades Norte, road to Bajo La Paz, ca. km 3, along the Río San Pedro, 10°08'58.7"N 84°34'03.3"W, 1300 m, premontane wet forest, secondary and remnants of primary vegetation, R.L. Dressler 7011, A. Karremans & F. Pupilin, 19 February 2008, flowered and pressed in cult. in Lankester Botanical Garden, July 2009, R.L. Dressler 7172 (CR). **Cartago:** Jiménez, Pejibaye, Alto del Humo, 23 May 2005, R.L. Dressler 6561, flowered and pressed in cult. in Lankester Botanical Garden, 14 August 2007, R.L. Dressler 6947 (CR). Paraíso, Orosi,




FIGURE 5. *Sobralia geminata* Dressler & Bogarín. A — Habit. B — Flower. C — Dissected perianth. D — Column and lip, lateral view. E — Column, frontal and side view. F — Pollinarium and anther cap. Drawing by D. Bogarín from the holotype.

Tapantí, road between Tapantí and Tausito, ca. 500 m before Tejos restaurant, 9°46'16.5"N 83°47'24.6"W, 1453 m, premontane wet forest, *R. Dressler 6661*, *M.A. Blanco & D. Bogarin*, 23 December 2005, flowered and pressed in cult. in Lankester Botanical Garden, 6 July 2006, *R.L. Dressler 6788* (CR). Turrialba, La Suiza, road between Pacayitas and La Suiza, 2 km south of Pacayitas, 9°52'29.9"N 83°35'03.6.4"W, 1150 m, premontane wet forest, 8 March 2006, *R.L. Dressler 6673*, *D. Bogarin*, *A. Karremans & F. Pupulin*, flowered and pressed in cult. in Lankester Botanical Garden, 10 July 2006, *R.L. Dressler 6789* (CR).

DISTRIBUTION: only known from Costa Rica.

ETYMOLOGY: from the Latin *geminatus*, “twinned,” referring to the usually paired flowers.

We first found small plants of this species in 2005. When they flowered, the flowers reminded us of *S. chrysostoma*, but we found the flowers to be consistently different from those of *S. chrysostoma*, so we thought it to be an unnamed species. In May of 2007 one of us (RLD) travelled to the northeast of Cartago Province, hoping to find a new species once found near Bonilla Arriba. There were colonies of a *Sobralia* along the roadsides and most of the youngest mature stems each had 2 capsules. My first thought was “The pollinators must be very efficient here,” but it was later clear that this population (Pacayas-Santa Cruz), at least, is self-pollinating. This was confirmed when other plants in the Lankester Botanical Garden often set fruits without pollination. Most of the plants produce only two flowers on each stem (either simultaneously or serially), so we have called it *Sobralia* “*geminata*,” because of the “twin” flowers. As one might expect from a partially self-pollinating species, there is local variation. Obviously, where the flowers are consistently selfed, bees do not carry mutant genes from one colony to another. In San Ramon, we found an attractive form that had the median groove on the lip a bit wider and sometimes had a slender, jointed, column appendage (in *Sobralia*, this appendage is more an “arm,” than a “wing”). This was so different from anything else we had seen in *Sobralia* that we thought it to be a new species. However, when the same plants flowered

again, they usually lacked the unique appendage, and they were self-pollinating.

The flowers of *Sobralia geminata* resemble those of *S. chrysostoma*, but they are consistently smaller, and they also have brownish spots or streaks on the lip, and some white near the apex. *Sobralia chrysostoma* frequently has dark purplish spots on the stems and sheaths, while the stems of *S. geminata* are consistently more heavily spotted than any other species in our area. The form of the stigma is quite variable in this species. The stigma varies a good deal in size, often being cuplike, as in most other species, or it may be straight, tongue-like, and subparallel with the column axis. In this latter pattern, the pollen probably germinates on the upper side of the stigma, thus accomplishing self-pollination.

Most *Sobralia* species are markedly synchronized (or “gregarious”) in their flowering, often flowering on the same day over a large area. One would not really expect a self-pollinating population to follow strict synchrony, yet the plants of this species in Lankester Botanical Garden usually flower on the same days. The plants collected south of Cartago have consistently flowered on the same days, even though they are quite self-pollinating. One doubts that there is much selective pressure for synchrony in self-pollinating plants, but, at the same time, there is probably little selective pressure against synchrony in such populations.

Our first impression of *S. geminata* was that it was predominantly self-pollinating. We have seen some selfing in most of the plants we have cultivated, as this description was in preparation, we had a number of plants flowering on 31 August 2009 and again on 9 September. Of the 16 flowers produced on 31 August, there were 6 capsules; while on 9 September there were 24 flowers, of which 15 remained as capsules. It is clear that the populations of Pacaya - Santa Cruz and Cartago - Cangreja are largely self-pollinating. One of the large plants from Cartago - Cangreja, however, had flowered heavily about 6 months earlier, and 3 capsules were allowed to remain on the plant. This plant has not produced any flowers in the recent flowerings. The two large plants from Cachi have flowered along with the rest of the *S. geminata*, but no capsules developed on either of these plants. Similarly, Ademar Silveira reports *S. geminata* in the buffer zone northeast of the




FIGURE 6. Distribution map of *Sobralia geminata* in Costa Rica.

Tapanti Refuge, but has observed no self-pollinating in the species in that area.

Now that we know *S. geminata*, we find plants of this species nearly everywhere we go in central Costa Rica. Indeed, we are rather tired of seeing them.

***Sobralia sotoana* Dressler & Bogarín, sp. nov.**

TYPE: PANAMA. Veraguas, collected near road north of Santa Fé de Veraguas, flowered in culture in Jardín Botánico Lankester, pressed from cultivation, 8 August 2009, R.L. Dressler 7168 (holotype: CR; isotype: JBL-spirit). FIG. 7, 8D.

Species *Sobralia aspera* Dressler & Pupulin similis, labello interius intense aurantiaco, callis basalibus farinosis differt.

Roots to 5 cm in diameter. *Stems* 33-75 cm, ca. 3 mm in diameter basally, 2 mm distally, sheaths slightly scurfy. *Leaves* elliptic or lance-elliptic, long acuminate, 15.0-23.5 x 2.2-5 cm. *Inflorescence* 3-5.0 x 0.5 cm, outer bract acuminate, inflorescence usually horizontal or nodding; subtending foliar bracts acuminate, 15-16 x 1.4-1.7 cm. *Ovary* sessile, 13-17 mm. *Sepaline tube* 12-15 mm. *Dorsal sepal* 3.3-4.5 x 1-1.6 cm, elliptic or lanceolate, subobtusate. *Lateral sepals* elliptic-lanceolate, tapering, 3.2-4.5 x 1.1-1.5 cm. *Petals* obovate, subobtusate, 3.1-3.5 x 1.2-1.3 cm.

Lip oblong-obovate, 4-4.2 x 2.8-3.1 cm, without keels; basal calli ca. 6 mm; column clavate, markedly thicker below stigma, 23-24 cm x 0.5 mm laterally x 4-4.5 mm dorsoventrally.

DISTRIBUTION: only known from Panama.

EPONYMY: In honor of Miguel Angel Soto Arenas, to whom this issue of Lankesteriana is dedicated.

When we visited Finca Dracula (Cerro Punta, Panama) in 2006, there were two relatively small *Sobralias* flowering. The staff at Finca Dracula referred to one as “Veraguas” and the other “Santa Fe,” but the city of Santa Fe is in Veraguas province, so these terms seemed a bit unclear. Erick Olmos explained that both species were found in the same area, near a lake north of Santa Fe, on the road that may eventually reach Calovébora, on the Caribbean coast. In any case, “Veraguas” proved to be indistinguishable from *Sobralia aspera*, earlier described from Costa Rica (Dressler & Pupulin 2008), while the other is clearly a distinct species. Further, the Costa Rican and the Panamanian plants of *S. aspera* flower on the same days, thus supporting their specific identity. Both of these resemble *Sobralia leucoxantha*, *S. macra*, *S. kruskayae* and *S. tricolor*, while *Sobralia* “Santa Fe” is clearly distinct from the other species just mentioned. The plant had produced very few flowers in the first year in Costa Rica, but in 2009 it has been quite floriferous.

Sobralia sotoana resembles *S. aspera* vegetatively, though the plant is a bit smaller. The upper stems are weakly but distinctly zig-zag, and the inflorescences much more markedly nodding, or reflexed, so the flowers are somewhat pendant. The flowers are a bit smaller and the lips are intense orange within with a narrow white margin.

ACKNOWLEDGEMENTS. We acknowledge the scientific services of the Costa Rican Ministry of Environment, Energy and Telecommunications (MINAET) and its National System of Conservation Areas (SINAC) together with Autoridad Nacional del Ambiente (ANAM) of Panama, for issuing the collecting permits under which wild species treated in this paper were collected, and for extending the necessary documentation to import orchid specimens from abroad for comparison. We are indebted to Andrés