

P. (§ Apodæ cæspitosæ) **HASTATA** sp. nov. *Caules secundarii graciles, ± 3 mm. longi. Folium circa 1.5 cm. longum, oblongi-lanceolatum, ad apicem tridentatum, carina prominente dorsali media. Pedunculus 2.5 cm. longus, gracilis, filiformis, circa 5-florus. Flores singulatim florescentes. Pedicelli 5 mm. longi, erecti, a rhachide contracta orti. Bracteæ inflorescentiæ circa 2 mm. longæ, approximatæ, scarriosæ, translucentes, oblongæ, ad apicem rotundatæ, abrupte in caudas aristiformes productæ. Sepala lateralia per longitudinem 2 mm. connata, oblongi-lanceolata, 7 mm. longa, 1 mm. lata, in caudas carnosas 2 mm. longas attenuata. Sepalum dorsale lateralibus angustius, simile. Petala linearia, 2.5 mm. longa, 1-nervia, sparsim serrata, acuta. Labellum unguiculatum, subhastatum, obtusum; lobus termina-*



PLEUROTHALLIS HASTATA

1, column; 2, petal; 3, labellum.

lis incrassatus et convexus ad apicem, 2 mm. longum, oblongum, margine dentato vel breviter fimbriato; lobi laterales triangulares, circa 1 mm. longi, subacuti, callo in angulo basali anteriore utroque, dente retrorso in latere utroque unguis. Columna gracilis, lamina membranacea circum clinandrium.

The flowers appear to have been whitish with purple spots, the labellum deep purple.

Alt. 350 m., Cubilquitz, Depart. Alta Verapaz, Guatemala, December, 1901, *H. von Tuerckheim* (no. 501).