

3. *Telipogon andicola*: aff. *T. tesselato* Lindl. (quam non vidimus) pedunculo folia acuta excedente, racemobi—plurifloro, bracteis triangulis carinatis, sepalis triangulis carinatis, tepalis cuneato obovatis obtuse acutis, labello subaequali. Folia sicca papyraceopergamenea lineariligulata sursum dilatata acuta, margine crispulo erosula duos prope pollices longa, duas tresve lineas lata, disticha, sicca superne favulosa, infimorum vaginæ a radicibus adventitiis laxis funiformibus multum flexuosis perforatae. Pedunculus anceps apice racemosus, bi—quinqueflorus. Bracteæ triangulæ argute carinatae pedicellis tripteris bene breviores. Ovarium tripterum demum capsulam ovoideam rostratam tripteram efficiens. Perigonium tenue flavulum nervis purpureis (in sicco). Sepala anguste triangula supra nervum medium alato carinata duas pollicis tertias longa, basi duas lineas lata. Tepala cuneato obovata obtuse acuta, sex—septemnervia nervulis evanidis nullisve: maculae violaceæ in ima basi papulis candidis introrsis hispida. Labellum subaequale apicem versus magis dilatatum obscure antice sinuato repandulum utrinque quadri—quinquenerve, nervulis extimis obliquis bene conspicuis — videtur obscure guttulatum fuisse; triangulum: in ima basi violaceo puberulum. Gynostemium humile bene hispidum atroviolaceum.

Western side of Pichincha on trees at 10000' of elev.
1848. 724.