


Xylobium Pickianum (Schltr.) L. Wms., *comb. nov.*
Bifrenaria Pickiana Schlechter in Orchis 6: 8, f. 1. 1912.

HONDURAS: flowers pale lilac, epiphyte in moist forest near Nacimiento del Río Lindo, Dept. Cortés, alt. 600 m., Aug. 6, 1948, *Williams & Molina 14555*; epiphyte in open mountain forest, Santa Cruz de Yojoa, Dept. Cortés, alt. 2000 feet, Sept. 7, 1933, *Edwards 541* (in Herb. Ames, Harvard Univ.).

COSTA RICA: region between Río Esquinas and Palmar Sur de Osa, vicinity of Esquinas Station, Prov. Puntarenas, alt. 30 m., July 17, 1950, *Allen 5584*; forested hills near Esquinas Station, vicinity of Río Esquinas, Prov. Puntarenas, alt. 60 m., Sept. 21, 1950, *Allen 5688*.

I had prepared a description of this plant, as a new species, when it discovered to be the same as Schlechter's *Bifrenaria Pickiana*, assumed to have been Colombian in origin.

The plate was prepared by Dorothy O. Allen from living material secured in Costa Rica.



XYLOBIUM PICKIANUM (Schltr.) L. Wms